

# The Variability of Active Galactic Nuclei and the Radial Transport of Vertical Magnetic Flux

Seok Jae Park

Korea Astronomy Observatory, San 36-1, Hwaam-Dong Yuseong-Gu, Daejeon 305-348,

Korea I: sjpark@apissa.issa.re.kr

and

Ethan T. Vishniac

Department of Astronomy, University of Texas, Austin, TX 78712

E:ethan@astro.as.utexas.edu

Received \_\_\_\_\_; accepted \_\_\_\_\_

## ABSTRACT

We consider the radial buoyancy of vertical magnetic field lines in radiation and gas pressure dominated accretion disks. We find that in addition to radial drift driven by turbulent diffusion and biased by the global field geometry, there are buoyancy effects which tend to move magnetic flux outward. In gas pressure dominated disks the poloidal magnetic field will move outward at a rate comparable to its inward advection. On the other hand, in a radiation pressure dominated disk the poloidal magnetic field will usually move outward faster than it is advected inward. This implies that the fields in disks in active galactic nuclei are generated at small radii by an internal disk dynamo. This conclusion can be avoided if the external field imposes a supersonic Alfvén speed within the disk without giving rise to interchange instabilities. In any case we note that variations in the mass transfer rate will lead directly to a modulation of the nonthermal emission from the disk system.

## 1. Introduction

It is well known that the emission-line spectra of active galactic nuclei (hereafter AGNs) are strongly variable. Typical time scales for this variation range continuously from the time scales appropriate for light travel across the inner regions of the accretion disk around a black hole, i.e.

$$t \approx GM/c^3, \quad (1)$$

(hours for supermassive black holes) up to years. This variability remains one of the central issues in any theoretical model of the central engine in an AGN.

A popular model for the source of the nonthermal radiation from an active galactic nucleus (AGN) consists of a supermassive black hole surrounded by a magnetized accretion disk. In its simplest form this model is taken to be axisymmetric and stationary. The nonthermal radiation is ascribed to the Blandford-Znajek process (Blandford and Znajek 1977), which emits nonthermal radiation from an extended magnetic field embedded in the black hole event horizon, the surrounding plasma, and the accretion disk. This model was reformulated and extended by Thorne and Macdonald (1982), Macdonald and Thorne (1982), and Thorne et al. (1986) in the ‘3+1’-spacetime formalism. The magnetic field in the disk will also give rise to a magnetically driven wind, and possibly a jet, as matter spirals out along the poloidal field lines (cf. Blandford & Payne 1982, Najita & Shu 1994, Shu, Najita, Ruden, & Lizano 1994, and Shu, Najita, Ostriker, Ruden, Wilken, & Lizano 1994). In addition, the ejection of toroidal flux from the disk may lead to winds and jets (e.g. Contopoulos 1995).

Starting from the axisymmetric, stationary model for emission due to the Blandford-Znajek process Park and Vishniac (1989*a*, 1989*b*) explored a time-dependent model which included the effects of variations in the mass accretion rate. The main point was to add the secular effects of mass accretion to the original axisymmetric, stationary model of

the nonthermal radiation. Such variations are expected because AGN accretion disks are expected to be radiation pressure and electron scattering dominated, and this can give rise to thermal and viscous instabilities on a broad range of time scales (Lightman & Eardley 1974). In these two papers we investigated the axisymmetric, nonstationary electrodynamics of a black hole and its accretion disk, respectively.

In the latter paper we found that the electrodynamic power output from the accretion disk can be variable on time scales associated with secular disk instabilities. The point was that the local fluctuations in fluid velocities in the accretion disk will cause fluctuations in the nonthermal component of the radiation by the Blandford-Znajek process. The time scales for these fluctuations, therefore, will reflect the range of orbital periods in the inner annulus of the disk.

In this paper we discuss another effect which may explain the fluctuations in the nonthermal emission, the relative motion of the infalling matter and the entrained poloidal field. The traditional view of the magnetic field in astrophysical objects, like accretion disks, is that the matter and magnetic field lines are tightly anchored to one another. Consequently accretion disks should continuously accrete poloidal field from the surrounding medium. Since the back reaction from the field on the matter flow will also increase without limit this leads to a situation in which the magnetic field completely overwhelms the gas pressure and drastically alters the properties of the accretion disk.

On the other hand, if one supposes that the magnetic field is not tightly coupled to the underlying fluid, then one can invoke the concept of turbulent diffusion for the magnetic field. This leads to the prediction that large scale magnetic fields tend to move outward in accretion disks and are eventually ejected from their outer edges (Van Ballegooijen 1989). Although it is difficult to reconcile turbulent diffusion with the view that the magnetic field fills the entire fluid volume, it is consistent with the idea that the magnetic field in stars

and accretion disks naturally forms intense, partially evacuated, fibrils that are capable of moving relative to the bulk of the fluid, i.e. flux tubes. In a recent paper Vishniac (1995a) examined the underlying physics behind the formation of such flux tubes and compared the predictions of a particular model with numerical simulations. A subsequent paper (Vishniac 1995b) predicted that the flux tubes in ionized accretion disks will be largely empty and discussed the consequences of this model for the vertical buoyancy of the toroidal field of an accretion disk. Here we extend this work to the radial motion of a poloidal field and show that in a radiation pressure dominated disk it leads to a strong outward motion of the field lines. This in turn leads to a modulation of the nonthermal power emitted by the magnetosphere as changes in the state of the disk raise and lower the rate at which the poloidal field is dispersed outward.

In §II of this paper we discuss the radial buoyancy of vertical magnetic field lines embedded in an accretion disk. In §III we discuss some of the implications of this work for nonthermal radiation from AGN.

## 2. The Radial Buoyancy of $B_z$

In order to understand the transport of flux within an accretion disk we need to have some model for the turbulent motions within the disk. Our work here is based on the idea that in an accretion disk these motions are driven by the magnetic field with the result that the angular momentum transport takes place within the disk and is mediated both by hydrodynamic transport and shearing of the local magnetic field. A magnetized accretion disk will be subject to a violent shearing instability which will act to transport angular momentum outward (Velikhov 1959, Chandrasekhar 1960, Balbus & Hawley 1991). The corresponding dimensionless viscosity  $\alpha$  will be roughly the ratio of magnetic to thermal

energies or

$$\alpha \sim \frac{B^2}{8\pi P} \sim \left( \frac{V_A}{c_s} \right)^2, \quad (2)$$

where  $B$  and  $P$  are the characteristic values of the magnetic field and pressure, respectively, in the disk, and  $V_A$  and  $c_s$  are the corresponding values of the Alfvén speed and sound speed. The resulting turbulence is anisotropic, but only by factors of order unity. This turbulence will be characterized by an eddy size of order  $V_A/\Omega$  and a typical turbulent velocity of order  $V_A$ . Larger scale modes, as well as the Parker instability, will be largely suppressed by these eddies (Vishniac & Diamond 1992). Nevertheless, there will be a residual mean vertical buoyant velocity of order  $\alpha c_s$ . A similar result applies when the magnetic field is considered to consist of individual flux tubes instead of some diffuse, space-filling mean field (Vishniac 1995b). For a thin disk the sound speed  $c_s \sim H\Omega$ , where  $H$  is the disk thickness and  $\Omega$  is the rotational frequency. Consequently, the dominant turbulent eddy size will be

$$L_T \sim \alpha^{1/2} H. \quad (3)$$

The magnetic field in a turbulent medium will tend to separate into regions of high and low magnetic field density. Most of the magnetic energy will be contained in flux tubes whose internal density is kept at a level well below that of the ambient medium through turbulent pumping (Vishniac 1995a). These flux tubes will interact with the surrounding fluid through the turbulent drag of the fluid as it moves by. Their mutual interactions guarantee that they will be broadly distributed in the fluid. The force per unit length on a flux tube is

$$\rho_t \frac{d^2 \vec{V}_t}{dt^2} (\pi r_t^2) = C_d \rho |\vec{V}_T - \vec{V}_t| (\vec{V}_T - \vec{V}_t) r_t + \hat{n}_c (\pi r_t^2) \frac{B_t^2}{4\pi R_c} - (\rho - \rho_t) \vec{g} (\pi r_t^2), \quad (4)$$

where the subscript  $t$  denotes quantities evaluated within the flux tubes,  $R_c$  is the radius of curvature of the local magnetic field,  $\vec{V}_T$  refers to the local fluid velocity,  $C_d$  is the coefficient of turbulent drag (of order unity),  $\vec{g}$  is the local gravitational acceleration, and  $\hat{n}_c$  is a unit

vector pointing in the direction of the curvature of the flux tube. We will subsequently use  $|V_T|$  to refer to the root mean square turbulent velocity in the fluid. Only the velocity components perpendicular to the local magnetic field enter into equation (4). If the flux tubes are largely empty, as they are in highly ionized disks, then the right hand side of this equation must sum to zero. Under most circumstances the buoyant force term will be much smaller than the other two terms (Vishniac 1995b). The properties of the flux tubes are largely independent of the resistivity of the gas provided that

$$\mathcal{M}_T^4 \left( \frac{|V_T|}{k_T \eta} \right) \left( \frac{4E_B}{f_c w E_T} \right)^{1/n+5/2} > \left( \frac{4}{C_d \gamma} \right)^2, \quad (5)$$

where  $\mathcal{M}_T$  is the turbulent Mach number,  $k_T$  is the wave number of the large scale eddies,  $\eta$  is the resistivity,  $E_B$  and  $E_T$  are, respectively, the spatially averaged magnetic and kinetic energy densities,  $f_c$ , and  $w$  are constants of order unity,  $\gamma$  is the adiabatic index of the fluid, and  $n$  is the turbulent power spectrum index (equal to 2/3 for Kolmogorov turbulence).

The viscosity can be ignored if

$$1 < \left( \frac{C_d \gamma}{4\pi^3} \right) \mathcal{M}_T^2 \left( \frac{|V_T|}{k_T \nu} \right) \left( \frac{4E_B}{f_c w E_T} \right)^{3/2+1/n}. \quad (6)$$

These conditions define the ideal fluid regime, within which the flux tubes are largely evacuated, except for a surface layer whose width is proportional to the square root of the resistivity. In accretion disks the kinetic and magnetic energy densities are roughly comparable due to the Balbus-Hawley instability. In this case, equations (5) and (6) can be rewritten as (Vishniac 1995b)

$$\alpha^3 \left( \frac{c_s H}{\eta} \right) \left( \frac{P}{P_{gas}} \right)^2 > \sim 16, \quad (7)$$

and

$$120 < \sim \alpha^2 \left( \frac{c_s H}{\nu} \right) \left( \frac{P}{P_{gas}} \right), \quad (8)$$

where  $P$  is the total pressure, and  $P_{gas}$  is the gas pressure. For temperatures and Mach numbers appropriate for ionized disks, both of these conditions are usually satisfied. In this

regime the typical flux tube radius is (Vishniac 1995a, Vishniac 1995b)

$$r_t \sim \alpha^{3/2} H \frac{P}{P_{gas}}. \quad (9)$$

The factor of  $P/P_{gas}$  enters here because turbulent pumping is ineffective for photons in accretion disks. Consequently, the magnetic pressure in a flux tube is limited by the ambient gas pressure instead of the total pressure.

In order to calculate the effects of radial buoyancy, we need to reconsider the third term in equation (4). The disk properties will vary radially, and the flux tube’s length and radius will change as it moves. We need to calculate the change in energy associated with moving a flux tube radially. If we consider an empty flux tube of radius  $r_t$  and length  $L$  embedded in an accretion disk with an ambient pressure  $P$  a length  $L$  then it will displace a volume of fluid equal to its own volume. Removing (adding) such a flux tube from an accretion disk will cause the surrounding gas to expand (contract) and lead to a drop (rise) in the local energy due to pressure work. The energy involved is

$$U_{tube} = \Delta \left[ \int_{-\infty}^{\infty} \left( \frac{1}{2} \rho z^2 \Omega^2 + \frac{1}{\gamma - 1} P \right) dz \right]. \quad (10)$$

If we assume that the gas expands adiabatically to fill the volume of the flux tube as it moves, then we can use the virial theorem for a pressure supported thin accretion disk, i.e.

$$\int_{-\infty}^{\infty} \rho z^2 \Omega^2 dz = \int_{-\infty}^{\infty} P dz, \quad (11)$$

to show that

$$U_{tube} = \int_{-\infty}^{\infty} P(z) \pi r_t^2(z) \frac{dL}{dz} dz \sim P(\pi r_t^2) L, \quad (12)$$

where we take the pressure and flux tube radius to be the appropriate values near the midplane of the disk. In other words, we obtain the usual expression for the energy loss due to adiabatic expansion. The contribution from the gravitational energy term, which is of comparable order, is canceled by an additional thermal energy term. This approach fails

for a flux tube in a radiation pressure dominated environment because only the gas pressure contributes to the energy expended by the gas expanding into the volume previously occupied by the flux tube. On the other hand, the gravitational potential term depends only on the matter density, and is not reduced by the tendency of the photons to fill up the otherwise empty flux tube. Consequently,  $U_{tube}$  is of order  $Pr_t^2 L$  rather than  $P_{gas} r_t^2 L$ .

Moving a flux tube radially implies a change in  $U_{tube}$  and a corresponding radial buoyancy. The radial force on a flux tube is equal to the radial derivative of  $U_{tube}$ . Since a given flux tube can split, or combine with other flux tubes, as it moves outward we need to constrain the radial variation of  $U_{tube}$  by using the conservation of magnetic flux, i.e.  $B_t r_t^2 \propto P_{gas}^{1/2} r_t^2$  is a constant of motion. Consequently, the radial force per unit length due to buoyancy is

$$F_r \approx -P(\pi r_t^2) \partial_r \ln(P P_{gas}^{-1/2} L). \quad (13)$$

Since the magnetic flux tube threads a turbulent disk, its actual length is  $\sim H(H/L_T) \sim H\alpha^{-1/2}$ . This implies that

$$F_r \approx -P(\pi r_t^2) \partial_r \frac{1}{2} \ln(P^2 H^2 P_{gas}^{-1} \alpha^{-1}). \quad (14)$$

The sign of this radial derivative is not obvious, and may not even be the same for all radii and at all times. For a gas pressure dominated disk we have  $P = P_{gas}$  and  $\dot{M} \approx \alpha P H \Omega^{-1}$ . Consequently,

$$\partial_r \frac{1}{2} \ln(P^2 H^2 P_{gas}^{-1}) = \partial_r \frac{1}{2} \ln(\dot{M} c_s \alpha^{-2}), \quad (15)$$

and the buoyant force will point outward in a stationary disk. On the other hand, near a thermal transition front, with  $\partial_r T$  and  $\partial_r \dot{M}$  negative, magnetic flux lines can be pulled inward.

For a radiation pressure dominated disk  $\dot{M} \approx \alpha P H \Omega^{-1}$ ,  $P_{gas} \propto \Sigma H^{-1} P^{1/4}$ , and

$H \propto \dot{M}$ , so that

$$\frac{P^2 H^2}{P_{gas} \alpha} \propto T^7 \dot{M}^3 \Omega. \quad (16)$$

For a stationary disk this increases inward, so that once again we obtain an outward buoyant force on the magnetic field lines.

If we assume that the curvature of a flux tube across the disk is negligible (more on that later) we obtain the mean radial motion due to buoyancy by equating the radial buoyant force given above with the turbulent drag due to a systematic outward motion of a flux tube. Averaging the turbulent drag on the flux tube over many eddies we find that

$$\langle |\vec{V}_T - \vec{V}_t| (\vec{V}_T - \vec{V}_t) \rangle \approx -\frac{3}{2} \langle |\vec{V}_T - \vec{V}_t| \rangle \vec{V}_b \sim -|V_T| \vec{V}_b, \quad (17)$$

where we have assumed that  $\langle \vec{V}_T \rangle = 0$  and  $\langle \vec{V}_t \rangle = \vec{V}_b$ . Dropping constants of order unity we obtain

$$\frac{F_r}{L} \approx \rho |V_T| V_r r_t \sim \frac{U_{tube}}{r}. \quad (18)$$

This implies that

$$V_r \sim \frac{P}{P_{gas}} \alpha \frac{c_s^2}{r \Omega}. \quad (19)$$

In other words, the radial buoyancy of the field lines gives rise to an outward drift which is of order the inward drift of the matter times  $P/P_{gas}$ . This same procedure (balancing buoyancy with turbulent drag) was used in Vishniac (1995b) to obtain a vertical buoyant velocity of  $\sim \alpha c_s$ . Note that we have ignored the specific angular momentum of the gas entrained on the field lines. This is justified as long as the flux tubes are largely empty, as they are in ionized accretion disks. In a disk dominated by gas pressure, the inward moving matter will act to drag the field lines inward, and the actual motion of the field lines will be determined by the detailed balance between these two effects. However, in a radiation pressure dominated environment the field lines will usually move outward. This conclusion will be affected by viscous and thermal instabilities in AGN disks, which may temporarily produce radial gradients which will pull the magnetic field lines inward.

When the flux tubes are not empty they will exchange matter, and consequently angular momentum, with their environment at the eddy turn over rate, which is  $\sim \Omega$ . Neglecting other effects which will tend to equalize the specific angular momentum of material inside and outside flux tubes, we see that this exchange rate implies that the average specific angular momentum of the flux tubes can differ from the surrounding gas by as much as

$$\Delta(r^2\Omega_{tube}) \sim \frac{V_r}{\Omega} \partial_r(r^2\Omega_{gas}). \quad (20)$$

Of course, there will be large fluctuations in this quantity, since the magnetic field will drive an instability by creating pockets of gas and field with specific angular momenta both larger and smaller than the local average. This systematic bias in the specific angular momentum of gas in the flux tubes will lead to a radial force of  $\sim -\rho_t\Omega V_r$ . Comparing this to the buoyant force per unit volume and using equation (19) we have

$$\frac{-\rho_t V_r \Omega}{(\rho - \rho_t) c_s^2 / r} \sim \frac{\rho_t}{\rho - \rho_t} \frac{P}{P_{gas}} \alpha. \quad (21)$$

Since the radiation pressure differential across a flux tube is negligibly small (Vishniac 1995b) the magnetic pressure in a tube must be balanced by a drop in the matter pressure. Therefore,

$$\alpha \sim \frac{\langle B^2 \rangle}{8\pi P} < \frac{B_t^2}{8\pi P} \sim \frac{\Delta P_{gas}}{P} \sim \frac{(\rho - \rho_t)}{\rho} \frac{P_{gas}}{P}. \quad (22)$$

From this we conclude that the ratio of the forces due to the specific angular momentum of material inside flux tubes to the buoyant forces will be less than  $\rho_t/\rho$ , which is less than one. Of course, this limit will be approached only when the flux tubes are barely distinguishable from the rest of the fluid, so that the volume average magnetic pressure is close to the magnetic pressure in the flux tubes. If we can neglect specific angular momentum, then the radial velocity due to buoyancy depends on the flux tube radius and internal magnetic field only in the combination  $B_t^2 r_t$ , which is always of order  $V_T^2 L_T$  for a magnetic field embedded in a turbulent medium and in equipartition with the turbulent energy (Vishniac 1995a).

We conclude that the results presented here should apply to all accretion disks except for those where the resistivity is so large as to nearly obliterate all flux tube structure.

Up to now we have assumed that the magnetic field outside the accretion disk crosses the disk with a negligible bending angle. This will not be the case if the external field is driving a magneto-centrifugal wind (e.g. Blandford & Payne 1982). In this case the motion of the field can be determined by vertical mixing through the disk. We can see this effect for a weak external field by using the usual mean field theory, and ignoring the division of the magnetic field in the disk into flux tubes. Our argument is equivalent to the one given by Van Ballegooijen (1989). The only conceptual difference is that we have justified the concept of turbulent diffusion by appealing to a specific physical model for the microscopic structure of the magnetic field. Starting from the induction equation

$$\partial_t \vec{B} = \vec{\nabla} \times \vec{v} \times \vec{B}, \quad (23)$$

we can define a transient field  $\vec{b}$  which is the response of the large scale magnetic field,  $\vec{B}_0$ , to small scale turbulent motions. We have

$$\partial_t \vec{b} = \vec{\nabla} \times \vec{v} \times \vec{B}_0. \quad (24)$$

Substituting this back into the right hand side of the induction equation we obtain

$$\partial_t \vec{B}_0 = \vec{\nabla} \times \langle \vec{v} \times (\int^t \vec{v} \cdot \vec{\nabla} B_0 dt) \rangle - \vec{\nabla} \times \langle \vec{v} \times (\int^t \vec{v} \cdot \vec{\nabla} \vec{B} dt) \rangle, \quad (25)$$

for incompressible turbulence. The first term in this equation is the source of the turbulent dynamo, if any, and the second term gives rise to diffusive effects. If we consider only the diffusive terms which affect the evolution of  $B_z$  then we get

$$\partial_t B_z = -\partial_r (\langle v_z^2 \tau \rangle \partial_z B_r) + \partial_r (\langle v_r^2 \tau \rangle \partial_r B_z) + \partial_r (\langle v_z v_r \tau \rangle (\partial_z B_z - \partial_r B_r)), \quad (26)$$

where  $\tau$  is the correlation time of the turbulence, in this case  $\sim \Omega^{-1}$ . The last term will be small, not only because the off diagonal terms of the averaged stress tensor will be small,

but because the extra radial derivative of  $B_r$  will make it of order  $H/r$  compared to the first term. (Since the divergence of the mean magnetic field vanishes, the two parts of the last term are equal to each other.) This implies an outward drift of the vertical field lines given by

$$V_r = \langle v_z^2 \tau \rangle \frac{\partial_z B_r}{B_z} \sim \alpha c_s \tan \theta, \quad (27)$$

where  $\theta$  is the bending angle of the magnetic field as it passes through the disk. Similar, and more precise, estimates can be found in paper by Lubow et al. (1994) and Reyes-Ruiz & Stepinski (1996). A different result can be obtained by assuming that the level of turbulent diffusivity in the disk is much less than that required to explain radial angular momentum transport. This is conceivable only if the Alfvén velocity in the disk is supersonic, thereby suppressing the Balbus-Hawley mechanism. If the magnetic field is driving a wind, then  $\theta$  will be positive and of order unity and this will dominate over the radial buoyancy of field lines for  $(P_{gas}/P) > (H/r)$ . On the other hand, if the field’s geometry is not directly affected by the disk, and has a curvature scale comparable to  $r$  then  $\theta$  will be of order  $H/r$  and the resulting radial drift velocity will be of the same order as the inward velocity of the matter. In this case radial buoyancy effects will dominate for radiation pressure dominated disks. Finally, in the event that the magnetic field lines bend *inward* as they cross the disk, this diffusive effect will move the vertical field lines inward.

We note that this derivation neglects the tension due to the bending of the large scale field as it crosses the disk and the possibility that the angular momentum in the disk is transported out into a magnetically driven wind instead of radially within the disk. Both assumptions are justified for a weak external field. A precise estimate of when this will run into problems must necessarily await a consensus on the degree to which an external field can produce a torque. Successful wind and jet models based on an external poloidal field (Blandford & Payne 1982, Najita & Shu 1994, Shu, Najita, Ruden, & Lizano 1994, and Shu, Najita, Ostriker, Ruden, Wilken, & Lizano 1994) give an equivalent ‘ $\alpha$ ’ based on an

external torque which can be as much as

$$\alpha_{ext} \sim \frac{B_{ext}^2}{P} \frac{r}{H}, \quad (28)$$

but these models assume that large radial bending angles near the disk are possible. As mentioned above, this is reasonable only for an induced magnetic pressure in the disk which is at least as large as the ambient thermal pressure. At smaller values the external field will induce turbulence, and transport, within the disk.

Without a definite model for the external torque, it is difficult to compare external and internal torques, but there are two points to consider with respect to the relationship between the internal turbulent viscosity and  $B_{ext}$ . First, the formation of flux tubes implies that the same amount of flux is compressed into a smaller volume, giving rise to a higher internal rms  $B_z$  than  $B_{ext}$ . If the flux tube formation process is very efficient, as would expect in an ionized disk, then

$$B_{z,internal} \sim B_{z,ext}^{1/2} (8\pi P_{gas})^{1/4}. \quad (29)$$

Moreover, the  $\alpha_{int}$  associated with an imposed vertical field is  $\sim (V_{Az}/c_s)$  rather than  $\sim (V_{Az}/c_s)^2$  (Vishniac & Diamond 1993). Consequently, we have

$$\alpha_{int} > \left( \frac{B_{z,ext}}{\sqrt{4\pi\rho}c_s} \right)^{1/2}. \quad (30)$$

Of course, this limit neglects any magnetic field generated within the disk itself, but such a field will only increase  $\alpha_{int}$ . We see that in spite of the potential advantage inherent in external torques due to their extended moment arm, reflected in the factor of  $r/H$  in equation (28), internal torques are likely to dominate for  $V_{Az} < c_s$ .

Finally, we note that the second term in equation (26) will tend to smooth out radial gradients in  $B_z$ . When the scale length for such gradients is of order  $r$ , then this will move vertical magnetic flux radially at a speed comparable to the accretion velocity. This would

prevent radial infall from building up the vertical magnetic field at small radii to arbitrarily high values even in the absence of radial buoyancy. In gas pressure dominated disks it also prevents radial buoyancy from expelling vertical flux from the disk, even in the absence of a disk dynamo. In a radiation pressure dominated disk this effect will be smaller, and will allow a positive  $\partial_r \ln B_z$  of order  $(P/P_{gas})r^{-1}$ .

We conclude that the strong magnetic fields required for nonthermal emission from the central parts of AGN disks must be either be maintained by dynamo processes at small radii, or dragged in through a magnetically dominated, and relatively nonturbulent, outer disk. In the latter case, the vertical field will adjust to changes in the disk at a the characteristic rate of

$$\tau_B^{-1} = \frac{V_r}{r} \sim \alpha \frac{H}{r} \Omega. \quad (31)$$

The former case will allow the field to adjust on a dynamo time scale, which will be (Vishniac 1995b)

$$\tau_{dynamo}^{-1} \sim \left( \frac{P}{P_{gas}} \right) \alpha \quad (32)$$

The nature of the dynamo at small radii is beyond the scope of this paper. Most work on disk dynamos has concentrated on the generation of toroidal fields, but the same processes will produce a poloidal field, albeit at a much reduced rate. In this case the field lines will tend to open up as differential rotation pumps energy into the disk magnetosphere, ultimately producing a large scale poloidal field.

### 3. Implications for the Variability of AGN

We have examined the radial buoyancy of poloidal field lines embedded in an accretion disk. In general the field lines will move outward through buoyant forces at a rate comparable to the inward drift caused by accretion. This leaves the question of their

global evolution unsettled. However, in radiation pressure dominated disks radial buoyancy dominates over accretion unless the magnetic field is strong enough to suppress turbulence. It follows that the global magnetic field embedded in accretion disks around AGN is likely to have its origin in dynamo processes taking place at small radii. In reaching this conclusion we have assumed that the angular momentum transport in these disks is mediated by magnetic field instabilities. Since the enhanced vertical buoyancy of magnetic field lines in such disks will tend to decrease the efficiency of this process (Vishniac 1995b) it seems possible that this assumption may be violated in the presence of hydrodynamic transport processes. As an example, we cite the possible role of internal waves, which normally give  $\alpha \sim (H/r)^2$  (Vishniac & Diamond 1989), a dimensionless viscosity which is unlikely to dominate in disks where the radiation pressure is negligible.

What does this imply about the variability of nonthermal radiation from AGNs? If the poloidal field is generated from within the disk, and moves outward through turbulent reconnection and radial buoyancy, then both its generation rate and its outward migration rate will depend on conditions within the disk. Consequently, variations in the state of the disk will lead to changes in the basic parameters of the magnetosphere, which will lead directly to variations in its nonthermal radiation. Variations in the mass accretion rate in the inner disk will arise from instabilities operating at larger radii and will reflect the thermal rates at those radii, roughly  $\alpha\Omega(r)$ . Since the time scales for disk variations will approach the light travel time across the magnetosphere for the inner regions of disks accreting near the Eddington limit it will be necessary to include time derivatives in the formulae for the structure of the magnetosphere.

As long as the poloidal field has only a weak effect on the mass flow within the disk, compared to internal processes driving angular momentum flow, the effects discussed here are incapable of driving disk instabilities. However, if the radiation pressure dominated

part of the disk is subject to local instabilities over its entire radius, then such instabilities will lead directly to variations in the structure of the magnetosphere and the nonthermal emission it drives. For example, thermal transition fronts will produce narrow annuli with a large local variations in  $\partial_r \dot{M}$ , driving the magnetic flux in the direction of the local mass flow. Since the disk instabilities will reflect the thermal and mass transport time scales for a wide range of radii, from the inner edge of the disk out to the radius where the radiation pressure of the disk loses its dominance, these time scales will also be present in the nonthermal emission. The upper limit to these time scales will depend on the details of the disk structure, but will clearly be much longer than orbital periods in the inner part of the disk. However, the shortest time scales for variations driven in this way will be comparable to the period given in Eq. (1).

This work has been supported in part by NASA grants NAGW-2418 and NAG5-2773 (ETV) and by the Basic Research Project 95-5200-002 of the Ministry of Science and Technology, Korea (SJP).

## REFERENCES

- Balbus, S., & Hawley, J.F. 1991, ApJ, 376, 214
- Blandford, R.D., & Payne, D.G. 1982, MNRAS, 199, 883
- Blandford, R.D., & Znajek, R.L. 1977, MNRAS, 179, 433
- Chandrasekhar, S. 1961, Hydrodynamic and Hydromagnetic Stability, Oxford University Press: London
- Contopoulos, J., ApJ, 450, 616
- Lightman, A.P. & Eardley, D.M. 1974, ApJ, 187, 897
- Lubow, S.H., Papaloizou, J.C.B., & Pringle, J.E. 1994, MNRAS, 267, 235
- Macdonald, D. A., & Thorne, K. S. 1982, MNRAS, 198, 345
- Najita, J.R., & Shu, F.H. 1994, ApJ, 429, 808
- Park, S.J., & Vishniac, E.T. 1989a, ApJ, 337, 78
- Park, S.J., & Vishniac, E.T. 1989b, ApJ, 347, 684
- Reyes-Ruiz, M., & Stepinski, T.F. 1996, ApJ, 459, 663
- Shu, F.H., Najita, J.R., Ruden, S.P., & Lizano, S. 1994, ApJ, 429, 797
- Shu, F.H., Najita, J.R., Ostriker, E.C., Ruden, S.P., Wilken, F., & Lizano, S. 1994, ApJ, 429, 781
- Thorne, K. S., & Macdonald, D. A. 1982, MNRAS, 198, 339
- Thorne, K. S., Price, R. H., & Macdonald, D. A. 1986, Black Holes: The Membrane Paradigm, (Yale University Press: New Haven)

- Van Ballegooijen, A.A. 1989, in *Accretion Disks and Magnetic Fields in Astrophysics*,  
(Kluwer: Dordrecht) p. 99
- Velikhov, E.P. 1959, *Soviet JETP*, 35, 1398
- Vishniac, E.T. 1995a, *ApJ*, 446, 724
- Vishniac, E.T. 1995b, *ApJ*, 451, 816
- Vishniac, E.T., & Diamond, P.H. 1989, *ApJ*, 347, 435
- Vishniac, E.T., & Diamond, P.H. 1992, *ApJ*, 398, 561
- Vishniac, E.T., & Diamond, P.H. 1993, in *Accretion Disks in Compact Stellar Systems*, ed.  
J.C. Wheeler (World Scientific Press: Singapore), p. 41